

Original article

Feasibility of tele-ophthalmology for screening for eye disease in remote communities

Li-Sheng Chen*, Ching-Yao Tsai^{††}, Tzeng-Ying Liu[§], Tao-Hsin Tung[‡],
Yueh-Hsia Chiu*, Chang-Chuan Chan^{**}, Der-Ming Liou* and
Tony Hsiu-Hsi Chen^{††}

*Institute of Public Health and Institute of Health Informatics and Decision Making, School of Medicine, National Yang-Ming University, Taipei; [†]Department of Ophthalmology, Taipei Municipal Chung-Hsin Hospital; [‡]Institute of Public Health, School of Medicine, National Yang-Ming University, Taipei; [§]Health Bureau of Lienkiang County, Matsu; ^{**}Institute of Occupational Medicine and Industrial Hygiene, College of Public Health, National Taiwan University; ^{††}Institute of Preventive Medicine, College of Public Health, National Taiwan University, Taipei, Taiwan

Summary

We assessed the feasibility of tele-ophthalmology in a remote location, Tungyin, an island 200 km from Taiwan, which has no ophthalmologist. Screening for eye diseases was carried out among residents aged 40 years or more. A total of 113 subjects, approximately 31% of the whole population, were enrolled in the screening programme. Images were transmitted (via ADSL) to a retinal specialist in Taiwan for diagnosis. The average processing time, excluding the time for copying files, was 6.4 s (SD 2.1) per subject. Transmission took 60–90 s for most of the images (83%). The average time required to make a diagnosis for each subject, including data entry, was approximately 34 s (SD 18). In screening for retinopathy, the detection rate with digital imaging (8.8%) was two times higher than with indirect ophthalmoscopy (4.4%). In 12% of cases macular degeneration was identified, and in 6% there were mild or moderate problems with the optic disc. Community-based screening for four categories of eye disease was successfully demonstrated using store-and-forward tele-ophthalmology.

Introduction

Tele-ophthalmology may be useful in remote areas where there are difficulties in accessing an ophthalmologist. For example, acute ophthalmological conditions in an Australian rural emergency department have been managed by teleconsultation with ophthalmologists working in a specialist eye clinic 900 km away¹. A pilot astigmatism screening study evaluated the transmission of digital images of children's pupil light reflexes to an ophthalmologist for diagnosis². Studies have tested the feasibility of tele-ophthalmology in the examination of patients with

glaucoma^{3–5} as well as for diabetic retinopathy screening^{6–9} (though not all of the latter found that it was feasible). Some studies have demonstrated the feasibility of using a portable fundus camera and a digital indirect ophthalmoscope for screening for common blinding eye diseases^{10,11}.

Despite this body of work, there is still a need to assess the feasibility of tele-ophthalmology for population-based screening for retinopathy and certain other eye diseases in remote areas, which typically have fewer than one ophthalmologist per 100,000 population. According to Lamminen *et al.*¹² the success of implementing tele-ophthalmology depends on technical aspects (e.g. appropriate facilities, the sending and reading of digital messages) as well as organizational and operational factors.

We have therefore investigated whether tele-ophthalmology is feasible for community-based

Accepted 5 September 2004

Correspondence: Professor Tony Hsiu-Hsi Chen, Graduate Institute of Preventive Medicine, College of Public Health, National Taiwan University, Room 207, 19 Hsuehshou Road, Taipei, Taiwan (Fax: +886 2 2358 7707; Email: stony@episerv.cph.ntu.edu.tw)

screening for retinopathy and other eye diseases in a remote area that has no ophthalmologist.

Methods

For the present study we selected a remote location, Tungyin, one of the Matsu islets, which is 33 km from mainland China and 200 km from Taiwan (Fig 1). In Tungyin there are many elderly people who require screening for retinopathy and other eye diseases, including macular oedema, optic disc problems and glaucoma. However, there is no ophthalmologist. There is ADSL communication between Taiwan and Tungyin. A tele-ophthalmology community screening programme for these four selected eye problems was tested.

Study population

There were 367 residents aged 40 years or more in Tungyin. The study population (i.e. those enrolled to attend the screening programme) comprised 113 residents, or approximately 31% of the target population. Their mean age was 53 years (SD 13). The most common reason for being unable to attend the screening programme among the 254 non-attenders was that they were not in Tungyin at the time, but were travelling (e.g. to visit relatives).

Only residents aged 40 years or older were invited to participate, but 14 residents aged 30–39 years also attended.

Study design and clinical assessment

The study design varied across the four different categories of eye disease of interest: retinopathy, macular oedema, optic disc problems and glaucoma. For the detection of retinopathy, we used both indirect ophthalmoscopy (one of the established tools for




Fig 1 Locations of Tungyin in the Matsu islands and Taiwan.

diabetic retinopathy screening) and digital imaging. We compared detection rates between digital imaging and indirect ophthalmoscopy. The detection of macular oedema and optic disc problems was entirely based on digital imaging. The diagnosis of glaucoma was made according to the anatomical findings from the patient's optic nerve disc, and functional visual field examination by frequency-doubling perimetry (FDP). Intraocular pressure (IOP) was also evaluated. An elevated IOP was defined as over 17 mmHg (1 mmHg=133 Pa). Severe glaucoma was defined as an optic cup:disc ratio over 0.7 with an FDP defect or elevated IOP. Mild glaucoma was defined as an optic cup:disc ratio between 0.7 and 0.5, or disc asymmetry of over 20%, with an FDP defect or elevated IOP.

The equipment and measures used in the clinical assessments included objective auto-refraction, measurements of presenting and best-corrected visual acuity, measurement of IOP with a non-contact tonometer, measurements of ocular dimensions, automated FDP, external and anterior segment examinations with a slit-lamp biomicroscope, and examinations of the vitreous, retina and optic nerve with an indirect ophthalmoscope. Digital 35-degree colour fundus images were obtained with a non-mydratric digital fundus camera (CR6-45, Canon, Japan).

Tele-ophthalmology procedure

The examination procedure is shown in Fig 2. Images of each fundus were checked using the non-mydratric




Fig 2 Procedure for the tele-ophthalmology screening programme.

digital fundus imaging system with two-field images. Images were transmitted to a retinal specialist in Taiwan for diagnosis. The retinal specialist reviewed each image, made any relevant diagnosis and made recommendations for subsequent care.

Each image was 2048 × 1360 pixels, at a colour depth of 24 bit/pixel. As previous studies had shown that JPEG compression of 1:28 for colour images was acceptable, fundus photographs were compressed 1:20 before transmission¹³. Transmission was by ADSL. The lowest upload speed was 64 kbit/s and the download speed was 512 kbit/s.

Results

Time required to process the image files

The first step was to transfer retinal images from the digital camera and to name each file appropriately. Two assistants compressed the image files. The average processing time, excluding the time for copying files, was 6.4 s (SD 2.1) per subject (Table 1).

Time required for image transmission

Assistants started to transmit the files using batch processing after every 20–30 subjects. Because of problems with the ADSL link in the remote area, it was

Table 1 Time required for renaming and compressing image files

Session number	Number of subjects	Time (s)	Mean time per subject (s)
1	24	180	7.5
2	32	240	7.5
3	7	60	8.6
4	32	180	5.6
5	18	60	3.3
<i>Total</i>	<i>113</i>	<i>720</i>	<i>6.4</i>

Table 2 Number of transmission attempts and time required

Number of transmission attempts required	Transmission time for image (s)						Total
	0–59	60–89	90–119	120–179	180–239	240+	
1	0	8	2	2	0	0	12
2	0	13	2	6	1	0	22
3	0	8	1	0	0	1	10
4	1	27	0	0	0	0	28
5	1	25	1	0	0	0	27
6	0	5	0	0	0	0	5
7	0	3	0	0	0	0	3
8	1	1	0	0	0	0	2
9	0	4	0	0	0	0	4
<i>Total</i>	<i>3 (3%)</i>	<i>94 (83%)</i>	<i>6 (5%)</i>	<i>8 (7%)</i>	<i>1 (1%)</i>	<i>1 (1%)</i>	<i>113</i>

sometimes necessary to attempt to transmit the images up to nine times. Table 2 shows the number of attempts and time required for the successful transmission of the image files. The majority of image files (88%) required five or fewer attempts at transmission. For most of the subjects (83%), transmission took 60–90 s (Table 2).

Time required for image diagnosis

The retinal specialists used a Web browser to view the digital images. They had been previously trained to use the application software, which was easy to use (it is available upon request from the first author). For diagnosis, each image could be zoomed in or out for further viewing. The average diagnosis time for each subject, including data entry, was approximately 34 s (SD 18).

Results of screening

Table 3 shows the retinopathy results for all 113 subjects. Retinopathy was graded according to a modified Airlie House Classification System¹⁴. Digital imaging detected 10 cases of 'other retinopathy' (six cases in the right eye and four cases in the left). By comparison, indirect ophthalmoscopy detected only five cases of 'other retinopathy' (three in the right eye, two in the left). No diabetic retinopathy was detected. Images from three subjects were undiagnosable due to failure of the digital fundus imaging system. In screening for retinopathy, the detection rate for digital imaging (8.8%) was two times higher than with indirect ophthalmoscopy (4.4%).

The screening results for the other three eye problems are summarized in Table 4. Among the 113 participants, we found 14 cases of early age-related macular degeneration (AMD) and found 14 other cases of AMD. In the optic disc assessment, we found six cases with mild problems and eight cases with moderate problems. There were no severe problems.

Table 3 Grade of retinopathy detected with retinal digital photography and indirect ophthalmoscopy

Retinopathy	Right eye				Left eye			
	Digital imaging		Indirect ophthalmoscopy		Digital imaging		Indirect ophthalmoscopy	
	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%
None	104	92	110	97	109	96	111	98
NPDR	0	0	0	0	0	0	0	0
Mild NPDR	0	0	0	0	0	0	0	0
Severe NPDR	0	0	0	0	0	0	0	0
PDR	0	0	0	0	0	0	0	0
Severe PDR	0	0	0	0	0	0	0	0
Other retinopathy	6	5	3	3	4	4	2	2
Insufficient image quality for diagnosis	3	3	0	0	0	0	0	0
<i>Total</i>	<i>113</i>	<i>100</i>	<i>113</i>	<i>100</i>	<i>113</i>	<i>100</i>	<i>113</i>	<i>100</i>

NPDR, non-proliferative diabetic retinopathy
PDR, proliferative diabetic retinopathy

Table 4 Screening results for macular degeneration, optic disc problems and suspected glaucoma

Eye disease	Right eye		Left eye	
	<i>n</i>	%	<i>n</i>	%
<i>Macular degeneration</i>				
None	102	90	96	85
Early age-related macular degeneration	6	5.3	8	7.1
Macular degeneration	5	4.4	9	8.0
<i>Optic disc assessment</i>				
Normal	105	93	107	95
Mild problems	3	2.7	3	2.7
Moderate problems	5	4.4	3	2.7
Severe problems	0	0	0	0
<i>Suspect glaucoma</i>				
None	96	85	96	85
Mild glaucoma	1	0.9	4	3.5
Severe glaucoma	2	1.8	2	1.8
High intraocular pressure	14	12.4	11	9.7
<i>Total</i>	<i>113</i>	<i>100</i>	<i>113</i>	<i>100</i>

The results of the comprehensive glaucoma assessment (IOP, IOP and fundus images) are also summarized in Table 4. We found five cases of mild glaucoma and four cases of severe glaucoma. High IOP (> 17 mmHg) (a risk factor for glaucoma) was found in 25 cases.

Discussion

Transmission methods in telemedicine can be grouped into three broad categories: store and forward, realtime and hybrid¹³. Store-and-forward systems relay data asynchronously. Information is acquired at one site, stored on a computer and then transmitted at a later time to another location, where it may be stored again

before review. Realtime systems work synchronously. Speakers, microphones and television cameras allow live videoconferences and group whiteboards to pass information almost instantaneously. Hybrid systems combine the capabilities of realtime and store-and-forward telemedicine¹³. Our study used store-and-forward methods to transmit image files because of their flexibility compared with realtime transmission. Although we had ADSL communication available between Taiwan and Tungyin, this was not always stable. Except when retransmission was required, the time required for renaming and compressing image files, the time required for image transmission, and the time required for diagnosis totalled approximately 2 minutes.

In the community screening study, no diabetic retinopathy was detected. This does not mean that our tele-ophthalmology method is inadequate for this purpose. The explanation is that because our target population was based on a general population, the prevalence diabetic retinopathy was low.

The results suggest that tele-ophthalmology can detect early age-related macular oedema (12% of the study population), optic disc problems (mild and moderate in a total of 12% of the study population) and high IOP (22%). Early treatment of these eye diseases can reduce the chance of blindness. However, the accuracy of detection through digital imaging requires further validation.

Two characteristics of the tele-ophthalmology screening programme deserve mention. First, our tele-ophthalmology was conducted from a mobile unit that moved between screening sites. Second, the target population was not patients who had sought medical service due to the presence of signs or symptoms but, unlike in earlier studies, was based on residents who participated annually in a multiple screening programme. This facilitated the early detection of eye disease.

Our study is not novel in the realm of telemedicine. However, the feasibility of a community-based tele-ophthalmology screening programme for the early detection of eye disease was demonstrated. It would be useful to know the sensitivity and specificity of the technique in the detection of posterior eye conditions. However, this would require all patients with negative findings to be sent to hospital for confirmatory diagnosis.

Cost is also an important factor in a screening programme. Our estimated cost per patient was about US\$10 (€8), which was slightly lower than that estimated in a study by Leese *et al.*¹⁵, who also used a mobile screening unit for diabetic retinopathy, in the UK. As our screening covered more than diabetic retinopathy, this suggests that our tele-ophthalmology was efficient.

Third, as suggested by Lamminen *et al.*¹², the quality of tele-ophthalmology is also important. Concerns have been raised as to whether lower image quality due to compression may affect diagnosis. In the present study this was unlikely, for two reasons. First, previous studies have shown that JPEG compression of 1:28 is not apparent to observers, and we chose a lower degree of compression, 1:20. Second, the digital imaging method had a higher detection rate for retinopathy than indirect ophthalmoscopy (see Table 3).

In conclusion, the community-based tele-ophthalmology screening programme was successful in remote Tungyin, Taiwan. The system was useful for the early detection of retinopathy, macular oedema, optic disc problems and glaucoma. Early treatment of these may reduce the incidence of blindness.

References

- 1 Rosengren D, Blackwell N, Kelly G, Lenton L, Glastonbury J. The use of telemedicine to treat ophthalmological emergencies in rural Australia. *Journal of Telemedicine and Telecare* 1998;**4** (suppl. 1):97-9
- 2 Researchers to test teleophthalmology on the reservation. *Telemedicine and Virtual Reality* 1997;**2**:121, 131
- 3 Yogesan K, Constable IJ, Fikelboom RJ, van Saarloos PP. Tele-ophthalmic screening using digital imaging devices. *Australian and New Zealand Journal of Ophthalmology* 1998;**26** (suppl. 1):9-11
- 4 Tuulonen A, Ohinmaa T, Alanko H, Hyytinen P, Juutinen A, Toppinen E. The application of teleophthalmology in examining patients with glaucoma: a pilot study. *Journal of Glaucoma* 1999;**8**:367-73
- 5 Lamminen H. Picture archiving and fundus imaging in a glaucoma clinic. *Journal of Telemedicine and Telecare* 2003;**9**:114-16
- 6 Williamson FH, Keating D. Telemedicine and computers in diabetic retinopathy screening. *British Journal of Ophthalmology* 1998;**82**:5-6
- 7 Schwartz SD, Harrison SA, Ferrone PJ, Trese MI. Telemedical evaluation and management of retinopathy of prematurity using a fiberoptic digital fundus camera. *Ophthalmology* 2000;**107**:25-8
- 8 Gomez-Ulla F, Fernandez MI, Gonzalez F, *et al.* Digital retinal images and teleophthalmology for detecting and grading diabetic retinopathy. *Diabetic Care* 2002;**25**:1384-9
- 9 Rotvold GI, Knarvik U, Johansen MA, Fossen K. Telemedicine screening for diabetic retinopathy: staff and patient satisfaction. *Journal of Telemedicine and Telecare* 2003;**9**:109-13
- 10 Schiffman JS, Li HK, Tang RA. Telemedicine enters eye care: practical experience. *Journal of Ophthalmic Nursing and Technology* 1998;**17**:102-6
- 11 Constable IJ, Yogesan K, Fikelboom R, Barry C, Cuypers M. Fred Hollows lecture: digital screening for eye disease. *Clinical and Experimental Ophthalmology* 2000;**28**:129-32
- 12 Lamminen H, Voipio V, Ruohonen K, Uusitalo H. Telemedicine in ophthalmology. *Acta Ophthalmologica Scandinavica* 2003;**81**:105-9
- 13 Li HK. Telemedicine and ophthalmology. *Survey of Ophthalmology* 1999;**44**:61-72
- 14 Klein R, Klein BE, Magli YL, *et al.* An alternative method of grading diabetic retinopathy. *Ophthalmology* 1986;**93**:1183-7
- 15 Leese GP, Ahmed S, Newton RW, *et al.* Use of mobile screening unit for diabetic retinopathy in rural and urban areas. *British Medical Journal* 1993;**306**:187-9